

CORRECTION

Open Access



Correction to: Dexketoprofen/tramadol 25 mg/75 mg: randomised double-blind trial in moderate-to-severe acute pain after abdominal hysterectomy

R. A. Moore^{1*}, H. J. McQuay², J. Tomaszewski³, G. Raba⁴, D. Tutunaru⁵, N. Lietuviene⁶, J. Galad⁷, L. Hagymasy⁸, D. Melka⁹, J. Kotarski¹⁰, T. Rechberger¹¹, B. Fülesdi¹², A. Nizzardo¹³, C. Guerrero-Bayón¹⁴, S. Cuadripani¹⁴, B. Pizà-Vallespir¹⁴ and M. Bertolotti¹³

Correction

Following publication of the original article [1], the authors reported that additional file 10 contained a typing error in the table “Percentage of responders ($\geq 50\%$ max TOTPAR) over two, four, six and eight hours (single-dose phase) (ITT Population)”. The table is to be read as follows:

		DKP/TRAM (N = 152)	DKP (N = 151)	TRAM (N = 150)	Placebo (N = 153)
TOTPAR ₂	Responder, n (%)	91 (60)	69 (46)	58 (39)	50 (33)
	Non-Responder, n (%)	60 (40)	82 (54)	91 (61)	103 (67)
Treatment comparisons <i>p</i> -value					
	DKP/TRAM vs. DKP	0.011			
	DKP/TRAM vs. TRAM	<0.001			
	DKP vs. Placebo	0.020			
	TRAM vs. Placebo	0.258			
TOTPAR ₄	Responder, n (%)	99 (65)	80 (53)	65 (43)	49 (32)
	Non-Responder, n (%)	52 (34)	71 (47)	84 (56)	104 (68)
Treatment comparisons <i>p</i> -value					
	DKP/TRAM vs. DKP	0.026			
	DKP/TRAM vs. TRAM	<0.001			
	DKP vs. Placebo	<0.001			
	TRAM vs. Placebo	0.038			

(Continued)

		DKP/TRAM (N = 152)	DKP (N = 151)	TRAM (N = 150)	Placebo (N = 153)
TOTPAR ₆	Responder, n (%)	105 (69)	72 (48)	64 (43)	49 (32)
	Non-Responder, n (%)	46 (30)	79 (52)	85 (57)	104 (68)
Treatment comparisons <i>p</i> -value					
	DKP/TRAM vs. DKP	<0.001			
	DKP/TRAM vs. TRAM	<0.001			
	DKP vs. Placebo	0.005			
	TRAM vs. Placebo	0.050			
TOTPAR ₈	Responder, n (%)	100 (66)	72 (48)	66 (44)	47 (31)
	Non-Responder, n (%)	51 (34)	79 (52)	83 (55)	106 (69)
Treatment comparisons <i>p</i> -value					
	DKP/TRAM vs. DKP	0.001			
	DKP/TRAM vs. TRAM	<0.001			
	DKP vs. Placebo	0.002			
	TRAM vs. Placebo	0.015			

TOTPAR: total pain relief; % max TOTPAR: percentage of the theoretical maximum possible TOTPAR; ITT: intention-to-treat; DKP/TRAM: dexketoprofen trometamol/tramadol hydrochloride 25 mg/75 mg; DKP: dexketoprofen trometamol 25 mg; TRAM: tramadol hydrochloride 100 mg; N: number of patients; n: number of patients with data. The ITT population included all patients randomised; response was defined as the achievement of at least 50% of the maximum

* Correspondence: andrew.moore@ndcn.ox.ac.uk

¹Pain Research & Nuffield Division of Anaesthetics, University of Oxford, The Churchill, Oxford, UK

Full list of author information is available at the end of the article



possible TOTPAR within the respective treatment arm; TOTPAR was calculated as the time-weighted sum of the pain relief (PAR) scores; PAR was measured on a five-point verbal rating scale (VRS) (0 = none, 1 = slight, 2 = moderate, 3 = good, 4 = complete); the percentage of PAR responders was analysed using a chi-square test.

Author details

¹Pain Research & Nuffield Division of Anaesthetics, University of Oxford, The Churchill, Oxford, UK. ²Balliol College, University of Oxford, Oxford, UK. ³Obstetrics-Gynaecology Private Clinic, Bialystok, Poland. ⁴Division of Gynaecology, Provincial Hospital in Przemysl, Przemysl, Poland. ⁵Genesys Fertility Center, Bucharest, Romania. ⁶Gynaecology, Riga East University Hospital Gynaecology Clinic, Riga, Latvia. ⁷GYNPOR, s.r.o, Sliac, Slovakia. ⁸Gynaecological Department, St. George Fejer County Teaching Hospital, Szekesfehervar, Hungary. ⁹Gynaecological Department, Latvian marine Medical Center, Riga, Latvia. ¹⁰Department of Gynaecological Oncology and Gynaecology, Medical University Hospital No 1, Lublin, Poland. ¹¹II Department of Gynaecology, Medical University Hospital No 4, Lublin, Poland. ¹²Department of Anaesthesiology and Intensive Care, University of Debrecen, Debrecen, Hungary. ¹³Clinical Research, Menarini Ricerche S.p.A – Menarini Group, Florence, Italy. ¹⁴Clinical Research, Laboratorios Menarini S.A. – Menarini Group, Badalona, Spain.

Received: 23 November 2017 Accepted: 23 November 2017

Published online: 30 November 2017

Reference

1. Moore RA, McQuay HJ, Tomaszewski J, Raba G, Tutunaru D, Lietuviene N, Galad J, Hagymasy L, Melka D, Kotarski J, Rechberger T, Fülesdi B, Nizzardo A, Guerrero-Bayón C, Cuadripani S, Pizà-Vallespir B, Bertolotti M. Dexketoprofen/tramadol 25 mg/75 mg: randomised double-blind trial in moderate-to-severe acute pain after abdominal hysterectomy. *BMC Anesthesiol.* 2016;16:9. <https://doi.org/10.1186/s12871-016-0174-5>.